

Resonant all-electric spin pumping with spin-orbit coupling

Valentina Brosco,^{1,2} Markus Jerger,³ Pablo San-Jose,^{2,4}
Gergely Zarand,^{5,6} Alexander Shnirman,^{7,6} and Gerd Schön^{2,6}

¹*Dipartimento di Fisica, Università “La Sapienza”, P.le A. Moro 2, 00185 Roma, Italy*

²*Institut für Theoretische Festkörperphysik, Karlsruhe Institute of Technology, 76128 Karlsruhe, Germany*

³*Physikalisches Institut, Karlsruhe Institute of Technology, 76128 Karlsruhe, Germany*

⁴*Instituto de Ciencia de Materiales de Madrid, CSIC,
Sor Juana Inés de la Cruz 3, E28049 Madrid, Spain*

⁵*Theoretical Physics Department, Institute of Physics,
Budapest University of Technology and Economy, Budapest, H-1521, Hungary*

⁶*DFG Center for Functional Nanostructures (CFN),
Karlsruhe Institute of Technology, 76128 Karlsruhe, Germany*

⁷*Institut für Theorie der Kondensierten Materie,
Karlsruhe Institute of Technology, 76128 Karlsruhe, Germany*

All-electric devices for the generation and filtering of spin currents are of crucial importance for spintronics experiments and applications. Here we consider a quantum dot between two metallic leads in the presence of spin-orbit coupling, and we analyze in the frame of a scattering matrix approach the conditions for generating spin currents in an adiabatically driven two-terminal device. We then focus on a dot with two resonant orbitals and show by specific examples that both spin filtering and pure spin current generation can be achieved. Finally, we discuss the effect of the Coulomb interaction.

PACS numbers:

Introduction. Many theoretical and experimental efforts have been devoted to spintronics, i.e., the design and control of spin-based electronic devices [1], and as a result it became possible to inject and filter spin polarized currents [2], to detect spin accumulation [3, 4, 5], and to produce ferromagnetic spin valves [6]. While an all-electrical control of spin currents would be clearly advantageous, so far most theoretical and experimental designs involve ferromagnetic leads or require the application of external magnetic fields.

Adiabatic pumping of charge in a cyclically modulated potential was first proposed by Thouless [7] and later studied in a variety of mesoscopic devices [8, 9, 10, 11, 12]. More recently, pumping of spins in nanostructures has been proposed as well, again with most mechanisms relying on external magnetic or exchange fields [13, 14, 15, 16, 17, 18], or the presence of ferromagnetic leads [19], and indeed in one experiment spin pumping in a magnetic field has been observed [20]. On the other hand, as first discussed in Refs. [16, 17], it is also possible to pump spin through quantum wires in the absence of external magnetic fields provided spin and orbit are coupled.

In the present paper we show that, in the presence of spin-orbit (SO) coupling, resonances associated with avoided level crossings of a quantum dot can be exploited to *pump* spin in a controlled way purely by cycling electrical gates. We shall focus on quantum dots with parameters such that the level spacing of the dot exceeds the typical width of the levels, $\delta\epsilon > \Gamma$. In this parameter regime individual states act as resonances, with position and coupling to the external leads which can be tuned by external gates [21]. *Resonant spin pumping* emerges,

when two of the levels lie close to the Fermi energy so that the SO coupling mixes them. We will show that in the vicinity of such resonant avoided level crossings the quantum dot can be used as an all-electric spin battery, and pumping cycles with transmitted spin of order $\hbar/2$ per cycle can be constructed. We also show that it is possible to choose the cycle parameters such that no charge is transferred through the quantum dot.

Scattering formalism. We consider an elastic scatterer coupled to two quasi-1D leads. In the left lead, far from the scattering region we can define longitudinal charge and spin current operators, J_L^0 and J_L^i , as

$$J_L^\mu(z, t) = -\frac{i}{2m} \int dx dy \left[\psi_L^\dagger \sigma^\mu (\partial_z \psi_L) - \text{h.c.} \right], \quad (1)$$

where σ^0 is the unit matrix, while σ^μ with $\mu = 1, 2, 3$ denote the usual Pauli matrices. The spinor field $\psi_L(x, y, z, t)$ destroys an electron in the left lead and ∂_z denotes the partial derivative with respect to the coordinate along the wire.

At low temperatures, we can use the approach of Brower [9, 17], to express the spin ($\vec{S}_L$) and charge (Q_L) pumped into the left lead within an adiabatic cycle as

$$Q_L = -\frac{e}{2\pi} \int_0^T \text{Im} \left[\text{Tr} \left\{ (\Lambda_L \otimes \sigma_0) \frac{d\mathcal{S}}{dt} \mathcal{S}^\dagger \right\} \right] dt, \quad (2a)$$

$$\vec{S}_L = -\frac{\hbar}{2\pi} \int_0^T \text{Im} \left[\text{Tr} \left\{ (\Lambda_L \otimes \vec{\sigma}) \frac{d\mathcal{S}}{dt} \mathcal{S}^\dagger \right\} \right] dt. \quad (2b)$$

Here $\mathcal{S} = \mathcal{S}(E_F, t)$ is the instantaneous scattering matrix at the Fermi energy, and Λ_L stands for the projector onto the left channel.

Following a strategy similar to Avron *et al.* [10], we decompose the scattering matrix to identify physical processes which contribute to spin and charge pumping. The presence of time-reversal symmetry implies that, in an appropriate basis, the scattering matrix is self-dual [23], i.e., $\mathcal{S} = \sigma_y \mathcal{S}^T \sigma_y$, and, for a quantum dot connecting two one-mode leads, $\mathcal{S}$ can be decomposed as

$$\mathcal{S} = U^0 U T U^\dagger U^0 \quad (3)$$

where the matrices U , U^0 and T are defined as follows:

$$\begin{aligned} U^0 &= \begin{pmatrix} e^{i\phi_L} & 0 \\ 0 & e^{i\phi_R} \end{pmatrix} \otimes \sigma_0, & U &= \begin{pmatrix} \mathcal{U}_L & 0 \\ 0 & \mathcal{U}_R \end{pmatrix}, \\ T &= \begin{pmatrix} -\sqrt{1-T_0} & \sqrt{T_0} \\ \sqrt{T_0} & \sqrt{1-T_0} \end{pmatrix} \otimes \sigma_0. \end{aligned} \quad (4)$$

where T_0 denotes the transmission coefficient of the dot and $\mathcal{U}_{L,R}$ are two-dimensional unitary matrices.

Substituting Eqs. (3-4) into Eqs. (2a-2b) we obtain

$$Q_L = \frac{e}{2\pi} \int_0^T \left[(1 - T_0) (\dot{\phi}_R - \dot{\phi}_L) \right] dt, \quad (5a)$$

$$\vec{S}_L = \frac{i\hbar}{2\pi} \int_0^T T_0 \text{Tr} \left\{ \left(\mathcal{U}_L^\dagger \vec{\sigma} \mathcal{U}_L \right) \left(\mathcal{U}_L^\dagger \dot{\mathcal{U}}_L - \mathcal{U}_R^\dagger \dot{\mathcal{U}}_R \right) \right\} dt. \quad (5b)$$

In Eq. (5a) we dropped a term $(e/2\pi) \int (\dot{\phi}_R + \dot{\phi}_L) dt$, representing the charge accumulated on the scatterer, which vanishes over a complete cycle. The first term in (5a) is finite even for $T_0 = 0$. This limit corresponds to peristaltic pumping, whereby a charge is first moved from the left lead to the dot, while the right contact is kept closed, and subsequently from the dot to the right lead with the left contact kept closed.

Interestingly, the pumped spin given by Eq. (5b) is proportional to T_0 . Therefore peristaltic spin pumping is not possible in the presence of time reversal symmetry. Also, while over a full cycle the charge is conserved, $Q_L + Q_R = 0$, the spin in general is not conserved: in particular if $[\mathcal{U}_L, \mathcal{U}_R] \neq 0$ we have $\vec{S}_R + \vec{S}_L \neq 0$. This is not surprising, since spin-orbit coupling – ultimately responsible for the spin pumping – allows for the transfer of angular momentum to the underlying lattice. We remark that the pumped spin in Eq. (5b) transforms as a vector under spin rotations in the left lead while it is invariant under spin rotation in the right lead, as it was also clear from the definition in Eq. (1).

Non-interacting quantum dot. To calculate explicitly the charge and spin transferred through a dot in the vicinity of an avoided level crossing, let us now introduce a non-interacting model to describe a dot with two orbital levels, $|n\sigma\rangle$ ($n = 1, 2$, $\sigma = \pm$) as shown in Fig.1. In the presence of (SO) coupling the isolated dot is described by the Hamiltonian $H^d = \sum d_{n\sigma}^\dagger \mathcal{H}_{n\sigma, n'\sigma'}^d d_{n'\sigma'}$,

with the operators $d_{n\sigma}^\dagger$ creating an electron in state $|n\sigma\rangle$ and the 4×4 matrix $\mathcal{H}^d$ given by

$$\mathcal{H}^d = \begin{pmatrix} \varepsilon_1 \sigma^0 & -i\vec{\alpha} \cdot \vec{\sigma} \\ i\vec{\alpha} \cdot \vec{\sigma} & \varepsilon_2 \sigma^0 \end{pmatrix}. \quad (6)$$

Here ε_n indicate the energies of the dot orbital states measured from the Fermi energy and the real vector $\vec{\alpha}$ is the SO field. Choosing the spin quantization axis parallel to $\vec{\alpha}$, we can express the full Hamiltonian as $H = \sum_\sigma H_\sigma$ with

$$\begin{aligned} H_\sigma &= \sum_n \varepsilon_n d_{n\sigma}^\dagger d_{n\sigma} + i \alpha s_\sigma \left(d_{1\sigma} d_{2\sigma}^\dagger - h.c. \right) \\ &+ \sum_{n, \xi, \lambda} \left(v_n^\lambda c_{\xi\sigma\lambda}^\dagger d_{n\sigma} + h.c. \right) + \sum_{\xi, \lambda} \xi c_{\xi\sigma\lambda}^\dagger c_{\xi\sigma\lambda}. \end{aligned} \quad (7)$$

with $s_\sigma = \pm 1$ for spin parallel/antiparallel to $\vec{\alpha}$ and $\alpha = |\vec{\alpha}|$. Here $c_{\xi\sigma\lambda}^\dagger$ creates a conduction electron of energy ξ and spin σ in lead λ . In the following, we shall assume that while the levels ε_n and the tunneling amplitudes v_n^λ can be tuned via gate voltages, $\vec{\alpha}$ remains approximately constant over a pumping cycle. Choosing $\vec{\alpha}$ as a quantization axis we can write the scattering matrix in spin-diagonal form, $\mathcal{S} = e^{i(\phi_L + \phi_R)} \mathcal{S}_\uparrow \oplus \mathcal{S}_\downarrow$, with $\mathcal{S}_\sigma$ parametrized using $\phi = \phi_L - \phi_R$ as

$$\mathcal{S}_\sigma = \begin{pmatrix} -e^{i\phi} \sqrt{1-T_0} & e^{is_\sigma\theta} \sqrt{T_0} \\ e^{-is_\sigma\theta} \sqrt{T_0} & e^{-i\phi} \sqrt{1-T_0} \end{pmatrix}. \quad (8)$$

Obviously, the phase ϕ is related to charge-pumping, while the phase θ determines the amount of spin pumped into the leads. If we only modify two external parameters, r_1 and r_2 , during a cycle, we can use Stokes theorem to recast the pumped charge and spin as

$$Q_L = -\frac{e}{2\pi} \iint d^2r B_c^{r_1 r_2}, \quad (9)$$

$$\vec{S}_L = \hat{\alpha} \frac{\hbar}{4\pi} \iint d^2r B_s^{r_1 r_2}, \quad (10)$$

where the charge and spin Brouwer's fields are defined as $B_c^{r_1 r_2} = \partial_{r_1} T_0 \partial_{r_2} \phi - \partial_{r_2} T_0 \partial_{r_1} \phi$ and $B_s^{r_1 r_2} = \partial_{r_1} T_0 \partial_{r_2} \theta - \partial_{r_2} T_0 \partial_{r_1} \theta$, respectively.




Figure 1: Orbital level structure of the quantum dot coupled to the leads. Due to the spin orbit field, $\vec{\alpha}_{so}$, pumping gives rise to charge and a spin current.

In the rest of the paper we shall study specific pumping cycles with two pumping parameters, either the tunnel couplings or the positions of the dot levels, appearing in the Hamiltonian H . To use Eqs. (9) and (10), we need to compute the transmission T_0 and the angles ϕ and θ in terms of these quantities. This can be done most easily by relating $\mathcal{S}$ to the Green's function G^d of the dot [26],

$$\mathcal{S}_{\sigma}^{\lambda\lambda'} = \delta_{\lambda\lambda'} - 2\pi i \sqrt{\rho_{\lambda}\rho_{\lambda'}} \sum_{n,n'} v_n^{\lambda} v_{n'}^{\lambda'*} G_{n,n'\sigma}^d(\varepsilon_F),$$

$$[G^d(\omega)]_{nn',\sigma}^{-1} = \omega \delta_{nn'} - (\mathcal{H}_{\sigma}^d)_{nn'} + i\pi \sum_{\lambda} \rho_{\lambda} v_n^{\lambda*} v_{n'}^{\lambda}.$$

where ρ_{λ} is the density of states in lead λ at the Fermi energy. From these equations we can obtain T_0 , ϕ , and θ in terms of the bare parameters of the dot. Introducing $w = v_{1L}v_{2R} - v_{2L}v_{1R}$ we can express for ϕ and θ as follows

$$\tan(\theta) = \frac{\alpha w}{\varepsilon_1 v_{2L}v_{2R} + \varepsilon_2 v_{1L}v_{1R}}, \quad (11)$$

$$\tan(\phi) = \pi\rho \frac{\varepsilon_1(v_{2L}^2 - v_{2R}^2) + \varepsilon_2(v_{1L}^2 - v_{1R}^2)}{\varepsilon_1\varepsilon_2 - \alpha^2 + \pi^2\rho^2w^2}. \quad (12)$$

Clearly, both levels are involved in spin pumping: if one of the levels is decoupled from the leads then $w = 0$, and only charge pumping is possible. Coupling the first and second level to the two leads with equal amplitudes also leads to a vanishing of the spin current.

In the following we analyze two kinds of cycles, “orbital energy cycles” and “tunnel coupling cycles”, where only the orbital energies of the levels are varied or also the tunnel coupling to the leads. Spin filtering and pure spin pumping can be achieved in both cycles.

Orbital energy cycles. For orbital energies cycles the relevant components of the Brouwer's fields are $B_s^{\varepsilon_1, \varepsilon_2}$ and $B_c^{\varepsilon_1, \varepsilon_2}$. For weak SO coupling these components have a dipolar structure in the plane $(\varepsilon_1, \varepsilon_2)$, as shown in the upper panels of Fig. 2. As for the single-level dot, studied in Ref. [24], the maxima of the charge field correspond to points of resonant transmission. The resonances of the spin field are instead located along the line $\varepsilon_1^r + \varepsilon_2^r = 0$ where the renormalized energies, ε_1^r and ε_2^r , are defined as $\varepsilon_1^r = \varepsilon_1 v_{2L}v_{2R}/w$ and $\varepsilon_2^r = \varepsilon_2 v_{1L}v_{1R}/w$. As we will show, the structure of the fields and the amount of pumped charge and spin depend very sensitively on the ratio between tunneling amplitudes and SO coupling.

To start with, let us consider the special case of a *symmetric dot* with a symmetric and an anti-symmetric orbital. In this case the Hamiltonian commutes with the operator $\pi \equiv \Sigma \otimes P$, where Σ and P denote the spin inversion and parity operators, and the tunneling amplitudes satisfy the following relations: $v_{1L} = v_{1R} = v_1$ and $v_{2L} = -v_{2R} = v_2$. In this symmetrical situation, a variation of the level energies, ε_1 and ε_2 , generates a *pure spin current* independently of the details of the cycle. In fact, as already noted by Aleiner *et al.* [27], parity

imposes additional constraints on the scattering matrix that along with time-reversal symmetry lead to the vanishing of Brouwer's charge field. In Fig. 2 (lower left panel) we plot the total spin pumped through a symmetric dot with $v_1 = v_2 = \sqrt{\Gamma/(2\pi\rho)}$ for a triangular pumping cycle such that $(\varepsilon_1 + \varepsilon_2) \in [\varepsilon_1 - \varepsilon_2, \infty]$ and $(\varepsilon_1 - \varepsilon_2) \in [-\infty, \infty]$. As one can see the pumped spin depends non-monotonously on the ratio α/Γ , the maximum occurring at $\alpha/\Gamma = 1$. Finally, in Fig. 2 (lower right panel) we plot the spin and charge pumped through a non-symmetric dot in a triangular cycle bounded by the lines, $|\varepsilon_1 - \varepsilon_2| \leq |\varepsilon_1 + \varepsilon_2|$ and $0 \leq |\varepsilon_1 + \varepsilon_2| \leq 10\Gamma$. In this case both the pumped charge and spin are non-vanishing and depend non-monotonously on the SO-coupling.

Tunnel coupling cycles. In the experiments it may be difficult to vary independently the energies of the two levels. For this reason we now consider pumping cycles where only the tunneling rates to the left lead and the offset of the two-levels with respect to the Fermi energy in the leads are varied. Specifically, we express the tunneling rates as follows, $v_{kM} = c_{kM} \sqrt{\Gamma_M/(2\pi\rho)}$ with $k \in [1, 2]$ and $M \in [L, R]$ and we assume that only Γ_L can be varied using some external gate. The factors c_{kM} , describing the overlap between the k -th dot's level and the scattering states in lead M , are assumed to be constant during the cycle. In Fig. 3 (left panel) we plot Brouwer's spin field $B_s^{\Gamma_L, \varepsilon_1 + \varepsilon_2}$. In this case, again, Brouwer's charge field is non-vanishing, however, with an appropriate tuning of the cycle shape, we can have pure spin currents. This is shown in Fig. 3 (right panel) where we plot the




Figure 2: Upper Panels: Brouwer's charge and spin field, $B_s^{\varepsilon_1, \varepsilon_2}$, in the $((\varepsilon_1 - \varepsilon_2), (\varepsilon_1 + \varepsilon_2))$ -plane for $v_{2L} = -v_{1R} = 2v_{2R} = 1.25v_{1L} = \sqrt{\Gamma/(2\pi\rho)}$, $\alpha/\Gamma = 0.25$. Lower Panels: pumped spin through a symmetric dot (left) and pumped spin and charge through a non-symmetric dot (right) as a function of the SO coupling α for a triangular cycle: $(\varepsilon_1 + \varepsilon_2) \in [0, 10\alpha]$ and $(\varepsilon_1 - \varepsilon_2) \in [-(\varepsilon_1 + \varepsilon_2), \varepsilon_1 + \varepsilon_2]$. In the symmetric dot case the tunnel couplings are $v_{2L} = -v_{1R} = v_{2R} = v_{1L} = \sqrt{\Gamma/(2\pi\rho)}$ while in the non-symmetric dot case they are the same as those of the upper panel.

total charge and spin pumped for a rectangular cycle, $\Gamma_L/\alpha \in [0, 5]$ and $(\varepsilon_1 + \varepsilon_2)/\alpha \in [0, x]$ as a function of x .

Interaction effects. To observe resonant spin pumping, the width of the two levels involved needs to be smaller or in the range of the level spacing. In this regime interaction effects, neglected so far, become important, and lead to a Coulomb blockade if all levels are far away from the resonances, and to the formation of a mixed valence state close to resonance. However, in all cases, the ground state of the dot is a Fermi liquid, and therefore low temperature spin pumping can be described in terms of a scattering matrix, just as for the non-interacting system. The only difference is that the parameters of this scattering matrix depend in a very non-trivial way on the bare model parameters. Determining the precise functional dependence of the scattering matrix on the gate voltages and the level positions is a cumbersome task beyond the scope of the present paper. The qualitative structure of the “phase diagram” and the spin pumping field is sketched in Fig. 4 for a dot with large Coulomb interaction. While the Coulomb interaction suppresses double occupancy for $\varepsilon_{1,2} < 0$, its effects are most likely not crucial in the regime, $\varepsilon_1, \varepsilon_2 > 0$, where the levels are only partially occupied. There non-interacting theory is expected to work reasonably well, and the resonant spin pumping discussed here should be observable.

Conclusions. We have studied spin and charge pumping in a quantum dot with spin-orbit interaction coupled to two metallic leads. Using Brouwer’s scattering approach to pumping, we first analyzed the general restrictions imposed by time-reversal symmetry on the pumped current. We then focused on the case of a dot with two resonant levels and showed that a spin of the order of $\hbar/2$ can be pumped in a cycle. We analyzed in particular two kinds of cycles involving the offset of the levels to the Fermi energy in the leads and, either the difference between the level energies or the tunneling to the leads. We demonstrated that, in both cases, with an appropriate tuning of the parameters, a controlled amount of spin




Figure 3: Left Panel: Brouwer’s spin field $B_s^{\Gamma_L, (\varepsilon_1 + \varepsilon_2)}$ in the $(\Gamma_L/\alpha, (\varepsilon_1 + \varepsilon_2)/\alpha)$ -plane for $\Gamma_{kM} = c_{kM}^2 \Gamma_M$ with $k = 1, 2$ and $M = L, R$, $c_{1L} = c_{2L} = 0.8$, $c_{1R} = 0.7$, and $c_{2R} = 0.6$. The two levels are assumed degenerate, $\varepsilon_1 - \varepsilon_2 = 0$, with zero hybridization, $\Delta = 0$, and the coupling to the right lead is $\Gamma_R/\alpha = 3$. Right Panel: Pumped spin and charge for a rectangular cycle $\Gamma_L/\alpha \in [0, 5]$ and $(\varepsilon_1 + \varepsilon_2)/\alpha \in [0, x]$ as a function of x .




Figure 4: Sketch of the effect of interactions.

and charge current can be pumped through the device. Finally we discussed the effect of Coulomb interaction, which should introduce only inessential corrections in the regime of partial dot occupation.

Acknowledgements- We were supported by the EU FP7 under Grant Agreement No. 238345 (GEOMDISS) and by the DFG-Schwerpunktprogramm 1285 ”Halbleiter-Spintronik”.

-
- [1] S. A. Wolf *et al.*, Science **294**, 1488 (2001).
 - [2] F. J. Jedema, A. T. Filip, and B. J. van Wees, Nature **410**, 345 (2001); K. Uchida *et al.*, Nature **455**, 778 (2008).
 - [3] R. M. Potok *et al.*, Phys. Rev. Lett. **89**, 266602 (2002).
 - [4] X. Lou, *et al.*, Nature Phys. **3**, 197 (2007); P. Kotissek *et al.*, Nature Physics **3**, 872 (2007).
 - [5] S. M. Frolov *et al.*, Nature **458**, 868 (2009).
 - [6] J. C. Sankey *et al.*, Nature Phys. **4**, 67 (2007).
 - [7] D. J. Thouless, Phys. Rev. B **27**, 6083 (1983).
 - [8] M. Büttiker, H. Thomas, and A. Prêtre, Z. Phys. B **94**, 133 (1994).
 - [9] P. W. Brouwer, Phys. Rev. B **58**, R10135 (1998).
 - [10] J. E. Avron *et al.*, Phys. Rev. B **62**, R10618 (2000).
 - [11] Y. Makhlin and A. D. Mirlin, Phys. Rev. Lett. **87**, 276803 (2001).
 - [12] M. Moskalets and M. Büttiker, Phys. Rev. B **68**, 075303 (2003).
 - [13] E. R. Mucciolo, C. Chamon, and C. M. Marcus, Phys. Rev. Lett. **91**, 258301 (2002).
 - [14] T. Aono, Phys. Rev. B **67**, 155303 (2003).
 - [15] R. Citro and F. Romeo, Phys. Rev. B **73**, 233304 (2006).
 - [16] M. Governale, F. Taddei, and R. Fazio, Phys. Rev. B **68**, 155324 (2003).
 - [17] P. Sharma and P. W. Brouwer, Phys. Rev. Lett. **91**, 166801 (2003).
 - [18] A. Schiller and A. Silva, Phys. Rev. B **77**, 045330 (2008).
 - [19] J. Splettstoesser, M. Governale, and J. König, Phys. Rev. B **77**, 195320 (2008).
 - [20] S. K. Watson *et al.*, Phys. Rev. Lett. **89**, 146802 (2003).
 - [21] K. MacLean *et al.*, Phys. Rev. Lett. **98**, 036802 (2007).
 - [22] D. R. Yafaev *Mathematical scattering theory*, (AMS, Providence, RI, 1992).
 - [23] C. W. J. Beenakker, Rev. Mod. Phys. **69**, 731 (1997).
 - [24] Y. Levinson, O. Entin-Wohlman, and P. Wölffe, Physica A **302**, 335 (2001).
 - [25] I. L. Aleiner and V. I. Falko, Phys. Rev. Lett. **87**, 256801 (2001).

- (2001).
- [26] Tai Kai Ng and P. A. Lee, Phys. Rev. Lett. **61**, 1768 (1988).
- [27] A. Aleiner, B.L. Altshuler, and A. Kamenev, Phys. Rev. B **62**, 10373 (2000).